arXiv:1303.3064v2 [cond-mat.stat-mech] 15 May 2013

Locality and thermalization in closed quantum systems

J. Sirker, N. P. Konstantinidis, and N. Sedlmayr
Department of Physics and Research Center OPTIMAS,
Technical University Kaiserslautern, D-67663 Kaiserslautern, Germany

(Dated: May 16, 2013)

We study the non-equilibrium dynamics of observables for finite one-dimensional quantum systems. By calculating the dependence of time and statistical averages on system size we investigate the role of the local and the exponentially many non-local conserved quantities for thermalization. We derive, in particular, a Mazur-type equality to split observables into a local and a non-local part with time and statistical averages for the local part being identical by construction. We then demonstrate that fluctuations in the non-local part are, in general, responsible for thermalization irrespective of whether or not the much stricter condition posed by the eigenstate thermalization hypothesis holds.

PACS numbers: 05.30.Ch,05.70.Ln,75.10.Pq

Introduction Starting a classical many-body system from a *typical* initial configuration we expect that the system equilibrates at long times so that *typical* observables become time independent. If the ensuing dynamics is ergodic, then the *ergodic theorem* [1–4] tells us that we can replace the time average of observables by an ensemble average. The ensemble provides a probability measure ρ on phase space which has to be invariant under time evolution because it describes the stationary state reached after the system has thermalized. The probability measure thus has to be a function of the conserved quantities $\mathcal{Q}_j$ with $\{\mathcal{H},\mathcal{Q}_j\}=0$ where $\{.,.\}$ is the Poisson bracket and $\mathcal{H}$ the Hamilton function of the system. In cases where $\mathcal{H}$ is the only independent conserved quantity this invariant phase space measure is the familiar microcanonical ensemble which becomes equivalent to the canonical one in the thermodynamic limit (TDL). Integrable systems with phase space dimension $2N$, on the other hand, have N independent conservation laws making them non-ergodic and restricting the motion in phase space to invariant tori. Importantly, the Kolmogorov-Arnold-Moser (KAM) theorem allows to describe the consequences of small integrability breaking perturbations on a quantitative level [4, 5].

Although recent theoretical [6–11] and experimental [12–15] studies have lead to new interesting insights, it is still fair to say that the non-equilibrium dynamics of a quantum system is much less well understood. One example, which has been studied numerically [8], is the expansion of a bosonic lattice gas into a larger empty lattice. In addition, also a number of other quantum quenches in lattice models with short-range interactions have been studied [9, 16–24]. In all cases the initial energy distribution becomes *singly-peaked* in the TDL with vanishing width [8]. Thermalization in these models has often been explained by the *eigenstate thermalization hypothesis* (ETH) [6–8, 25] which assumes that the expectation values of the considered operator in the eigenstate basis fluctuate little between eigenstates close in energy and can thus be directly replaced by an ensemble average.

In this letter we show that thermalization, in principle, can also occur in a more generic scenario where fluctuations between eigenstate expectation values close in energy are large. We derive a criterion for these fluctuations which has to be fulfilled if the system thermalizes.

We restrict our discussion to time-independent Hamiltonians H with short-range interactions in position space and a discrete spectrum. Position space, however, is by no means special: For a Hamiltonian with e.g. local interactions in momentum space the same approach can be used and the results directly carry over. The time average of an observable O for an initial pure state $|\Psi_0\rangle$ is defined by

$$\bar{O}=\lim_{\tau\to\infty}\frac{1}{\tau}\int_0^\tau dt\langle\Psi_0|\mathrm{e}^{iHt}O\mathrm{e}^{-iHt}|\Psi_0\rangle. \tag{1}$$

By using a spectral representation of the observable O and assuming that the spectrum of H is non-degenerate [45] we immediately obtain

$$\bar{O}=\langle O\rangle_{\mathrm{diag}}\;\equiv\;\langle\Psi_0|O_{\mathrm{diag}}|\Psi_0\rangle=\sum_n O_{nn}\underbrace{|\langle\Psi_0|n\rangle|^2}_{\equiv|c(n)|^2} \tag{2}$$

where $O_{\mathrm{diag}}=\sum_n O_{nn}P_n$. Here $O_{nn}=\langle n|O|n\rangle$ with $P_n=|n\rangle\langle n|$ being the projection operator onto the eigenstate $|n\rangle$ and $H|n\rangle=\varepsilon_n|n\rangle$. Off-diagonal elements of O do not contribute in time or statistical averages, i.e., to address thermalization it is sufficient to consider the diagonal part.

If we want to replace the time average by an ensemble average following the prescription in the classical case, we immediately encounter an important aspect which is different from the classical problem: All the projection operators are conserved, $[H,P_n]=0$, and the number of conserved quantities thus always equals the Hilbert space dimension D. A density matrix ρ_{diag} which yields a statistical average equal to the time average (2) thus apparently has to be a function of all the projection operators [26], e.g.,

$$\rho_{\text{diag}} = \exp(-\sum_n \lambda_n P_n)/Z_{\text{diag}} \tag{3}$$

with $Z_{\text{diag}} = \text{Tr}\,\rho_{\text{diag}}$ and Lagrange multipliers λ_n fulfilling the condition $e^{-\lambda_n}/Z_{\text{diag}} = |\langle\Psi_0|n\rangle|^2$. While it is easy to see that this ensemble indeed fulfills $\text{Tr}(O\rho_{\text{diag}}) = \langle O\rangle_{\text{diag}} = \bar{O}$, see Eq.(2), by construction and thus naively proves the ergodic theorem in the quantum case, it depends on the initial state and fixes the *microstate* of the system up to phase factors which are irrelevant for the long-time dynamics. In quantum statistical mechanics we are, however, only interested in the *macrostate* consisting of many microstates which cannot be distinguished by local probes [26].

The new important ingredient, which has to be taken into account in the quantum case, is the distinction between *local* and *non-local* conserved quantities. A local operator for a lattice model is defined as $\mathcal{Q}_n = \sum_j q_j^n$ where q_j^n acts on lattice sites $j, j+1, \cdots, j+n$ only, with n finite. In a field theory this then becomes a fully local operator $\mathcal{Q}_n = \int d\boldsymbol{r} q^n(\boldsymbol{r})$. In the TDL we expect that only the local conserved quantities are needed in the density matrix to describe the thermalized state. This number is usually finite for a generic quantum system while it increases linearly—but not exponentially!—with system size for an integrable one-dimensional model [27, 28]. Experiments on cold atomic gases as well as most numerical studies of the thermalization problem are, however, done on finite systems [8, 19, 29] where a distinction between local and non-local conservation laws, strictly speaking, does not exist. Understanding the scaling with system size of the local and non-local contributions to the ensemble is therefore a problem of practical relevance.

General considerations: A Mazur-type equality To obtain a condition for thermalization on the microscopic level of individual eigenstates we write

$$\begin{aligned} 0 &= \lim_{N\to\infty} \Delta O_N \equiv \lim_{N\to\infty} [\langle O\rangle_{\text{diag}} - \langle O\rangle_{\text{th}}] \qquad (4)\\ &= \lim_{N\to\infty} \sum_{n=1}^{D} \underbrace{\frac{\langle OP_n\rangle_{\text{th}}}{\langle P_n^2\rangle_{\text{th}}}}_{O_{nn}} \big(\underbrace{\langle\Psi_0|P_n|\Psi_0\rangle}_{|c(n)|^2} - \underbrace{\langle P_n\rangle_{\text{th}}}_{\rho_{\text{th}}^{nn}} \big) \\ &\overset{N\to\infty}{\to} \int d\varepsilon\; O(\varepsilon) [\underbrace{|c(\varepsilon)|^2\nu(\varepsilon)}_{\Gamma_{\text{diag}}(\varepsilon)} - \underbrace{\rho_{\text{th}}(\varepsilon)\nu(\varepsilon)}_{\Gamma_{\text{th}}(\varepsilon)}] \end{aligned}$$

where $\langle O\rangle_{\text{th}} = \text{Tr}\{O\rho_{\text{th}}\}$ with $\rho_{\text{th}} = \exp(-\sum_n \lambda_n \mathcal{Q}_n)/Z$, $Z = \text{Tr}\,\rho_{\text{th}}$ being the appropriate thermal density matrix including all the local conserved quantities $\mathcal{Q}_n$ with Lagrange parameters $\{\lambda_n\}$ determined by the set of equations $\langle\Psi_0|\mathcal{Q}_n|\Psi_0\rangle = \text{Tr}\{\mathcal{Q}_n\rho_{\text{th}}\}$ [21]. Furthermore, we have rewritten the matrix elements O_{nn} in the generic form of a projection. In the last line we have introduced a coarse-grained description in the thermodynamic limit with $O(\varepsilon) = \sum_{\varepsilon-d\varepsilon<\varepsilon_n<\varepsilon-d\varepsilon} O_{nn}$ and similarly a thermal energy distribution $\Gamma_{\text{th}}(\varepsilon)$ and initial energy distribution $\Gamma_{\text{diag}}(\varepsilon)$ both including the coarse-grained density of states $\nu(\varepsilon)$. The assumption of the ETH scenario is that O_{nn} becomes a smooth function of energy ε_n in the TDL, i.e., $O(\varepsilon = \epsilon_n) = O_{nn}$ so that a coarse graining is not required. One of the main points of our study is that this is unnecessarily restrictive. Note that both the initial energy distribution $\Gamma_{\text{diag}}(\varepsilon)$ as well as the thermal distribution $\Gamma_{\text{th}}(\varepsilon)$ will become sharply peaked in the TDL so that only a small energy window will contribute to the integral in the last line of Eq. (4). In order to investigate thermalization in more detail we explain in the following how to separate an observable into a part proportional to the local conserved quantities and an orthogonal part.

The distinction between local and non-local conserved quantities has turned out to be essential for quantum transport. Here it is known that a local or quasi-local [30] conserved quantity can protect a current from decaying while a non-local conserved quantity cannot [31–34]. This is understood by using a Mazur equality [31, 35, 36] where the current operator is written as a sum of projections onto all the conserved quantities of the system. In exactly the same way, we can project the considered observable onto the local and non-local conserved quantities to study thermalization. Note that this is nothing but a simple basis change in operator space. Instead of using $P_n = |n\rangle\langle n|$ as the basis we use the f many local conserved quantities which can be written as

$$\bar{P}_n \equiv \mathcal{Q}_n = \sum_{j=1}^{D} a_j^n P_j \;(n = 0, \cdots, f-1) \tag{5}$$

with coefficients a_j^n plus $D-f$ many non-local operators $\bar{P}_n$, $n = f, \cdots, D-1$ such that $\langle\bar{P}_n\bar{P}_m\rangle_{\text{th}} = \langle\bar{P}_n^2\rangle_{\text{th}}\delta_{nm}$ are orthogonal. Here we have choosen $\langle\cdots\rangle_{\text{th}} = \text{Tr}\{\cdots\rho_{\text{th}}\}$ as the natural inner product in the vector space of Hermitian operators.

To be concrete we consider the case that the system has only a single local conservation law—the Hamiltonian itself which can be written as $H = \sum_j \varepsilon_j P_j$. The thermal density matrix $\rho_{\text{th}} = \exp(-H/T)/Z$ is then just the usual canonical ensemble expected to describe the system at long times in the TDL. As a first step we choose the set of normalized operators $\{\tilde{H}, \tilde{P}_1, \tilde{P}_2, \cdots, \tilde{P}_{D-1}\}$ where $\tilde{H} = H/\sqrt{\langle H^2\rangle}$, $\tilde{P}_n = P_n/\sqrt{\langle P_n^2\rangle}$ with $P_n^2 = P_n$. We now define the normalized operator

$$\tilde{U} = (\tilde{H} - \tilde{P}_D)/\sqrt{\langle(\tilde{H} - \tilde{P}_D)^2\rangle} \tag{6}$$

so that

$$R(\tilde{H}) = \tilde{H} - 2\tilde{U}\langle\tilde{U}\tilde{H}\rangle = \tilde{P}_D \tag{7}$$

where $\tilde{P}_D$ is the projection operator which we have replaced by $\tilde{H}$. Applying the same orthogonal transformation (a so-called Householder rotation) onto the projection operators

$$R(\tilde{P}_i) = \tilde{P}_i - 2\tilde{U}\langle\tilde{U}\tilde{P}_i\rangle = \bar{P}_i \tag{8}$$

for $i = 1, 2, \cdots, D-1$ thus generates the required orthonormal set $\{\bar{P}_0 \equiv \tilde{H}, \bar{P}_1, \bar{P}_2, \cdots, \bar{P}_{D-1}\}$. With the help of this new basis in operator space we can split $O_{\rm diag}$ into a local and a non-local part

$$O_{\rm diag} = \underbrace{\frac{\langle OH\rangle_{\rm th}}{\langle H^2\rangle_{\rm th}} H}_{O_{\rm loc}} + \underbrace{\sum_{n=1}^{D-1} \frac{\langle O\bar{P}_n\rangle_{\rm th}}{\langle \bar{P}_n^2\rangle_{\rm th}} \bar{P}_n}_{O_{\rm nonloc}} . \tag{9}$$

Energy conservation during time evolution demands $\langle\Psi_0|H|\Psi_0\rangle = \langle H\rangle_{\rm th}$ which fixes the temperature T [8, 10, 17] and guarantees the equivalence of the diagonal and canonical ensemble average for the first term, $O_{\rm loc}$ in (9), which is proportional to H. Eq. (4) can thus be rewritten as a condition for the non-local part only.

We are typically interested in observables O, e.g. correlation functions, which are not affected by an energy shift $H' = H - E_0$. In this case we can simplify the condition on the non-local part further by demanding that $\langle O_{\rm nonloc}\rangle_{\rm th} = 0$. For this gauge we have to choose the reference energy

$$E_0 = \frac{\langle OH\rangle_{\rm th}\langle H\rangle_{\rm th} - \langle O\rangle_{\rm th}\langle H^2\rangle_{\rm th}}{\langle OH\rangle_{\rm th} - \langle O\rangle_{\rm th}\langle H\rangle_{\rm th}} . \tag{10}$$

The condition for thermalization (4) then reads

$$\begin{aligned} 0 &= \lim_{N\to\infty} \langle O_{\rm nonloc}\rangle_{\rm diag} = \lim_{N\to\infty} \underbrace{\sum_{n=1}^{D-1} \Delta\bar{O}_N(n)}_{\Delta O_N} \\ &= \lim_{N\to\infty} \sum_{n=1}^{D-1} \underbrace{\frac{\langle O\bar{P}_n\rangle_{\rm th}}{\langle \bar{P}_n^2\rangle_{\rm th}}}_{\bar{O}_{nn}} \langle\Psi_0|\bar{P}_n|\Psi_0\rangle \end{aligned} \tag{11}$$

where the last line defines the diagonal matrix elements $\bar{O}_{nn}$ in the new operator space basis. Using this new basis we have achieved a separation into a part $O_{\rm loc}$ with $\langle O_{\rm loc}\rangle_{\rm diag} \equiv \langle O_{\rm loc}\rangle_{\rm th}$ by construction and a part, $O_{\rm nonloc}$, for which the agreement between time and statistical average is non-trivial.

The model So far, our discussion has been general. To test the concepts we have introduced above, we want to study in the following a specific lattice model, the one-dimensional anisotropic Heisenberg model

$$\begin{aligned} H(\Delta, J_2) &= J\sum_{j=1}^{N} h_{j,j+1} + J_2 \sum_{j=1}^{N} h_{j,j+2}, \\ h_{i,j} &= \frac{1}{2}\left(S_i^+ S_j^- + h.c.\right) + \Delta S_i^z S_j^z, \end{aligned} \tag{12}$$

where S is a spin-1/2 operator, J (J_2) are the superexchange couplings for the nearest (next-nearest) neighbors, respectively, and Δ parametrizes an exchange anisotropy. In the following we set $J = 1$, use periodic boundary conditions, and study the model by exact diagonalization taking spin and spatial symmetries into account [37]. The model (12) is integrable for $J_2 = 0$. In this case the number of conserved local operators $\mathcal{Q}_n$ increases linearly with system size N. In the non-integrable case $J_2 \neq 0$, H itself and $S^z_{\rm tot} = \sum_j S^z_j$ are the only conserved local operators. To make contact with previous thermalization studies, we consider a quench scenario [38, 39]. As initial state we choose the ground state $|\Psi_0(\Delta, J_2)\rangle$ of the Hamiltonian (12) with parameters Δ and J_2. We then time evolve with $H(\tilde{\Delta}, \tilde{J}_2)$ where $(\tilde{\Delta}, \tilde{J}_2) \neq (\Delta, J_2)$.



FIG. 1: (Color online) (a) Quench with $|\Psi_0(5, 0.2)\rangle$ and $H(1, 0.2)$. Expectation values $\langle \boldsymbol{S}_i \boldsymbol{S}_{i+1}\rangle$ using different ensembles: (1) Difference between canonical and diagonal ensemble and linear fit, (2) between canonical and extended ensemble $\rho_{\bar{P}_1}$ and exponential fit, (3) between $\rho_{\bar{P}_1}$ and the diagonal ensemble and the difference of the two fits. (b) Difference of canonical and diagonal ensemble average for an integrable quench with $|\Psi_0(5, 0)\rangle$ and $H(1/2, 0)$.

Locality and statistical ensemble We start by studying the role played by non-local conserved quantities for the statistical ensemble. To do so we define an extended canonical ensemble

$$\rho_{\bar{P}_j} = \exp(-\beta H + \lambda_j \bar{P}_j)/Z, \tag{13}$$

where the two parameters β, λ_j are determined by $\langle\Psi_0|H|\Psi_0\rangle = {\rm Tr}\{H\rho_{\bar{P}_j}\}$ and $\langle\Psi_0|\bar{P}_j|\Psi_0\rangle = {\rm Tr}\{\bar{P}_j\rho_{\bar{P}_j}\}$. This extended ensemble is compared with the canonical and the diagonal ensemble in Fig. 1 for the generic case $j = 1$ with $\bar{P}_1$ as defined in Eq. (8). We have found that the qualitative results for the finite size scaling are independent of which of the $\bar{P}_j$ we choose in the extended ensemble (13) and consistent with the following: (a) The change in the average of an observable caused by including an additional non-local conservation law vanishes exponentially with system size. (b) The contribution of all non-local conservation quantities to the ensemble average vanishes linearly in $1/N$. This follows from the fact that the canonical ensemble is obtained from the diagonal one by neglecting all non-local quantities, see Eq. (3). For a finite system the non-local conserved quantities do, however, contribute showing that the TDL is essential for a full thermalization. In the integrable case shown in the inset of Fig. 1, the canonical ensemble does not describe



FIG. 2: (Color online) Longer-range correlations for the same quench as in Fig. 1(a). Shown is the scaling of the relative deviation between ensemble averages Δ_{rel} for (a) $|i-j| = 1, 2, 3, 4, 5$ (from bottom to top) and (b) for $|i-j| = N/2$.

the equilibrated state at long times because other *local* conserved quantities are not taken into account [40].

Locality and observables Next, we want to study how the amount of locality of the operator itself affects its thermalization. Numerical data for $O = \langle \boldsymbol{S}_i \boldsymbol{S}_j \rangle$, which is a two-point correlation function of a fully local operator, are shown in Fig. 2. The relative deviation $\Delta_{\mathrm{rel}} = |(\langle \boldsymbol{S}_i \boldsymbol{S}_j \rangle_{\mathrm{diag}} - \langle \boldsymbol{S}_i \boldsymbol{S}_j \rangle_{\mathrm{can}})/\langle \boldsymbol{S}_i \boldsymbol{S}_j \rangle_{\mathrm{diag}}|$ between the long-time mean (diagonal ensemble) and the canonical ensemble average becomes larger the larger the distance is and thus the less local O is, see Fig. 2(a). If we fix the distance between the spin operators to $N/2$ as in Fig. 2(b) then the canonical ensemble average approaches zero with increasing N much faster than the long-time mean so that $\Delta_{\mathrm{rel}} \to 1$. Thus not only the initial distribution $\langle \Psi_0 | \bar{P}_n | \Psi_0 \rangle \approx \langle \Psi_0 | P_n | \Psi_0 \rangle$, see Suppl. Mat., but also the operator-dependent first term in Eq. (11), $\bar{O}_{nn}$, has to play a crucial role in the thermalization process.

Projection onto locally conserved quantities To study the role of locality of the operator in more detail we now use the splitting into a local and a non-local part defined in Eq. (9). The diagonal distribution function $|c(\varepsilon_n)|^2$, and the microcanonical and canonical ones, $\Gamma_{\mathrm{mic/can}}(\varepsilon)$, for the same quench as before are shown in Fig. 3(a). While the diagonal distribution is expected to become singly peaked in the thermodynamic limit [8], this is clearly not the case for the system sizes within reach of exact diagonalization. The initial state distribution thus cannot be simply replaced by the microcanonical ensemble. Furthermore, O_{nn}, shown in Fig. 3(b1) and (b2) for two different correlation functions, shows large fluctuations. Thus ETH is not directly applicable, yet we already see clear indications for thermalization as shown in Fig. 1 and Fig. 2.



FIG. 3: (Color online) Same quench as in Fig. 1(a) for $N = 16$. Shown are: (a) the initial, microcanonical and canonical energy distribution functions, and results for $O = \boldsymbol{S}_i \boldsymbol{S}_{i+1}$ in (b) and $O = \boldsymbol{S}_i \boldsymbol{S}_{i+4}$ in (c). (b1) and (c1) show O_{nn} while (b2) and (c2) show $\bar{O}_{nn}$.

To understand these findings we return to the more generic condition for thermalization, Eq. (11), which only requires that the sum of the fluctuations in the non-local part averages out to zero. Indeed, $\bar{O}_{nn}$, shown in Figs. 3(b2) and (c2), has much smaller fluctuations centered around zero and both correlation functions display similar behavior once the local part is separated off. Although we see clear evidence for thermalization, no conclusive finite size scaling for the fluctuations in O_{nn} could be obtained so that the question whether or not the ETH holds in the considered case has to remain open. For $\bar{O}_{nn}$, however, we do obtain a clear finite size scaling (see Suppl. Mat.) showing that $\bar{O}_{nn} \to 0$ for all n in the TDL. Thus every single term in the sum in Eq. (11) becomes zero so that the projections of the observable onto the nonlocal conserved operators can be ignored allowing the system to thermalize to the canonical ensemble.

Conclusions To summarize, we have looked at the question of thermalization in closed quantum systems from the perspective of conservation laws. Contrary to a classical system two distinct kinds appear: local and nonlocal ones. By choosing a new basis in operator space we have separated the considered observable into a projection onto the locally conserved quantities, for which the time and statistical average agree by construction, and a projection onto the non-local ones where thermalization takes place. The condition for thermalization which is obtained by this separation involves fluctuations in the non-local part only and is more general than ETH. This, in principle, opens up the possibility that systems exist which do thermalize although the ETH does not apply.

The authors thank F.H.L. Essler, B. Fine, A. Polkovnikov, and M. Rigol for discussions and acknowledge support by the Collaborative Research Centre SFB/TR49 and the Graduate School of Excellence MAINZ.

Supplementary material

The purpose of our supplementary material is twofold. In Sec. we will first provide some additional information about the basis rotation required to split an observable into a local and a non-local part, numerical data testing the ETH which proved to be inconclusive, as well as some technical details about the coarse graining procedure used to obtain a continuous density of states. Secondly, we also provide some additional information about what happens in the case of integrable systems in Sec. .

Details of the calculation

In Sec. we give some details which are helpful to understand the splitting of an observable into a local and a non-local part in a more general context. In Sec. we present a finite size scaling analysis of the fluctuations in O_{nn} and $\bar{O}_{nn}$. Finally, we explain in Sec. the coarse graining procedures used to obtain the continuous distributions.

Splitting into local and non-local part

In the paper we have argued that the proper basis in operator space to study thermalization consists of the local conserved operators plus an additional set of non-local operators which span the rest of the operator space. In this basis the time and statistical average for the part of the observable obtained by projecting onto the local conserved operators agree by construction so that thermalization becomes a statement about the projection onto the non-local conserved operators only.

Choice of inner product As in Eq. (9) of the paper we can write the diagonal part of an arbitrary observable O for the case that H is the only local conserved operator as

$$O_{\text{diag}} = \underbrace{\frac{\langle OH \rangle}{\langle H^2 \rangle} H}_{O_{\text{loc}}} + \underbrace{\sum_{n=1}^{D-1} \frac{\langle O\bar{P}_n \rangle}{\langle \bar{P}_n^2 \rangle} \bar{P}_n}_{O_{\text{nonloc}}} \qquad (14)$$

with $\langle H\bar{P}_n \rangle = 0$ and $\langle \bar{P}_n \bar{P}_m \rangle = \delta_{nm} \langle \bar{P}_n^2 \rangle$. In the paper we have chosen $\langle \cdots \rangle = \langle \cdots \rangle_{\text{th}} \equiv \text{Tr}\{\cdots \rho_{\text{th}}\}$ as inner product in the vector space of Hermitian operators. While this is the natural choice, it is also possible to use any other proper inner product in operator space. The splitting into a local and a non-local part thus depends on the choice of the metric. This is, of course, expected since the very notion of distances requires the definition of a metric first.

Using a general inner product and projection operators $\bar{P}_n$ which are orthogonal with respect to this metric, Eq. (11) of the paper can be written in the more general form

$$\begin{aligned} 0 &= \lim_{N\to\infty} \Delta O_N \equiv \lim_{N\to\infty} [\langle O \rangle_{\text{diag}} - \langle O \rangle_{\text{th}}] \qquad (15) \\ &= \lim_{N\to\infty} \sum_{n=1}^{D-1} \underbrace{\frac{\langle O\bar{P}_n \rangle}{\langle \bar{P}_n^2 \rangle}}_{\bar{O}_{nn}} \big(\langle \Psi_0 | \bar{P}_n | \Psi_0 \rangle - \langle \bar{P}_n \rangle_{\text{th}} \big). \end{aligned}$$

We want to stress that ΔO_N does not depend on the choice of the inner product. It is only once we start considering the contributions on a microscopic level that this choice becomes relevant.

Energy shift Apart from the inner product there is another 'gauge degree of freedom' in the second line of Eq. (15). By shifting the energy, $H' = H - E_0$, the projection operators $\bar{P}_n$ are modified because of the orthogonality condition $\langle H\bar{P}_n \rangle = 0$. The qualitative behavior of $\bar{O}_{nn}$ is, however, not affected. A convenient, unique gauge is obtained by demanding that $\sum \bar{O}_{nn} \langle \bar{P}_n \rangle_{\text{th}} = 0$. For a general metric $\langle \cdots \rangle$ this is achieved by choosing

$$E_0 = \frac{\langle OH \rangle \langle H \rangle - \langle O \rangle \langle H^2 \rangle}{\langle OH \rangle + \langle O \rangle \langle H \rangle_{th} - 2\langle O \rangle_{th} \langle H \rangle}. \qquad (16)$$

for a given Hamiltonian H and operator O. By setting $\langle \cdots \rangle = \text{Tr}\{\cdots \rho_{\text{th}}\}$ this expression simplifies to Eq. (10) of the paper.

Relation between the original and the rotated basis Once an inner product has been chosen, the new basis consisting of the f local conserved quantities and $D - f$ non-local operators can be orthonormalized by using the Householder reflection given in the paper. To be concrete, we choose again $\langle \cdots \rangle = \text{Tr}\{\cdots \rho_{\text{th}}\}$ as the inner product and assume that H is the only local conserved quantity. The relation between the old and new operator basis can then be expressed as

$$\bar{P}_m = \sum_{n=1}^{D} a_n^m P_n = \sum_{n=1}^{D} \underbrace{a_n^m \sqrt{\langle P_n \rangle_{\text{th}}}}_{\langle \bar{P}_m \tilde{P}_n \rangle_{\text{th}}} \tilde{P}_n \ (m = 1, \cdots, D-1) \qquad (17)$$

where $\tilde{P}_n = P_n / \sqrt{\langle P_n^2 \rangle_{\text{th}}}$ with $P_n^2 = P_n$ are the normalized projection operators, $\langle \tilde{P}_i \tilde{P}_j \rangle_{\text{th}} = \delta_{ij}$. Using the definition of the Householder reflection and some simple algebra we find

$$\langle \bar{P}_m \tilde{P}_n \rangle_{\text{th}} = \delta_{nm} - \frac{2\varepsilon_n \varepsilon_m}{\langle (\tilde{H} - \tilde{P}_D)^2 \rangle_{\text{th}}} \sqrt{\langle P_n \rangle_{\text{th}}} \sqrt{\langle P_m \rangle_{\text{th}}}. \qquad (18)$$

While $\langle (\tilde{H} - \tilde{P}_D)^2 \rangle_{\text{th}} \propto \mathcal{O}(1)$ we have $\sqrt{\langle P_n \rangle_{\text{th}}} \propto \mathrm{e}^{-N/2}$ so that the expansion coefficients a_n^m are sharply peaked at $n = m$. As a consequence, the initial distribution is not affected by the rotation in the thermodynamic limit, i.e.,

$$\langle \Psi_0 | \bar{P}_n | \Psi_0 \rangle \overset{N\to\infty}{\to} \langle \Psi_0 | P_n | \Psi_0 \rangle \quad (n = 1, \cdots, D-1) \qquad (19)$$

and becomes sharply peaked in the thermodynamic limit. The matrix elements of the observable

$$\begin{aligned}\bar{O}_{mm} &= \frac{\langle O\bar{P}_m\rangle_{\rm th}}{\langle \bar{P}_m^2\rangle_{\rm th}} = \sum_n O_{nn}\langle P_n\bar{P}_m\rangle_{\rm th} \\ &= \sum_n a_n^m \langle P_n\rangle_{\rm th} O_{nn} \end{aligned} \tag{20}$$

are, however, changed because they are given by summing over the exponentially many old matrix elements O_{nn} so that the exponentially small corrections in a_n^m, see Eq. (18), still matter.

Fluctuations in O_{nn} and $\bar{O}_{nn}$ and the eigenstate thermalization hypothesis

According to the eigenstate thermalization hypothesis, O_{nn} should become a smooth function of the eigenenergy ε_n in the thermodynamic limit. For the system sizes we can exactly diagonalize we are clearly far from that limit and fluctuations in O_{nn} are large. Nevertheless, we can check how these fluctuations scale with system size.

We define the average size of the fluctuations in an interval:

$$\Delta_O = \sum_n{}' \left|O_{nn} - \langle O_{nn}\rangle_{mc}\right| . \tag{21}$$

The prime on the sum refers to a restriction to an energy interval of 0.05 times the bandwidth $W_\Delta = \varepsilon_D - \varepsilon_1$, and centered on the middle of the spectrum, $E = W_\Delta/2 + \varepsilon_1$, with ε_1 the ground state energy. This can be more precisely defined as

$$\begin{aligned}\sum_n{}' \cdots &= \sum_{n=1}^{D} \cdots \Gamma(\varepsilon_n - E), \text{ where} \\ \Gamma(\varepsilon) &= \theta\left[\varepsilon + 0.05W_\Delta\right] - \theta\left[\varepsilon - 0.05W_\Delta\right] \end{aligned} \tag{22}$$

and $\theta(\varepsilon)$ is the Heaviside function. $\langle O_{nn}\rangle_{mc}$ is the locally defined average, in other words the microcanonical ensemble, calculated here with the same energy window:

$$\langle O_{nn}\rangle_{mc} = \sum_{m=1}^{D} O_{mm}\Gamma(\varepsilon_m - \varepsilon_n) . \tag{23}$$

To compare the size of the fluctuations with the magnitude of the operator we define $\langle O\rangle_E = \sum_n' O_{nn}$. We can define the same in the rotated basis:

$$\Delta_{\bar{O}} = \sum_n{}' \left|\bar{O}_{nn} - \langle \bar{O}_{nn}\rangle_{mc}\right| \tag{24}$$

with the interval for the sum defined as for Eq. (21). Strictly speaking $\langle \bar{O}_{nn}\rangle_{mc}$ is no longer the microcanonical ensemble average as n no longer labels the eigenenergies. Nonetheless one can define an analogue and we retain the same notation for ease of presentation.



FIG. 4: (Color online) Quench with $|\Psi_0(5, 0.2)\rangle$ and $H(1, 0.2)$. Shown is a comparison of (a) $\Delta_{\bar{O}}$, (b) Δ_O, and (c) $\Delta_O/\langle O\rangle_E$ for different system sizes from $N = 8$ to 16. Plotted are the observables $O = \boldsymbol{S}_i\boldsymbol{S}_j$ with $|i-j| = 1$ (black circles), $|i-j| = 2$ (red squares), $|i-j| = 3$ (green triangles), $|i-j| = 4$ (blue diamonds), and $|i-j| = 5$ (purple triangles). $\Delta_{\bar{O}}$ shows clear exponential scaling to zero (note the logarithmic scales).

We consider again the same quench as in the paper with $|\Psi_0(5, 0.2)\rangle$ and $H(1, 0.2)$, and look at observables $O = \boldsymbol{S}_i\boldsymbol{S}_j$ for different $|i-j|$. In Fig. 4 we plot $\Delta_{\bar{O}}$ and Δ_O for system sizes $N = 8$ to 16. The absolute magnitude of the fluctuations of ΔO , Fig.4(b), can be several orders of magnitude larger than their average value even for $N = 16$, see Fig. 4(c), so that ETH can not yet be said to apply. In particular it does not seem possible that this can be the explanation for the data of Fig. 2(a) of the paper. Though the trend for the fluctuations appears to be for them to become on the whole smaller for larger system sizes, no clear-cut scaling can be seen. Furthermore, the relative size of the fluctuations, defined as $\Delta_O/\langle O\rangle_E$ as in Ref. [41], also shows no conclusive scaling with the system size, see Fig. 4(c). However, for the non-local part $\Delta_{\bar{O}}$, shown in Fig. 4(a), one sees clear scaling to zero which depends exponentially on the system size, supporting the division into local and non-local operators which we have implemented.

Energy distributions and coarse graining

In order to plot the continuum energy distributions a coarse graining is necessary. The density of states is first made continuous by approximating

$$\nu(\varepsilon) \equiv \sum_n \delta(\varepsilon - \varepsilon_n) \approx \sum_n \chi_W(\varepsilon - \varepsilon_n) , \tag{25}$$

with an envelope function:



FIG. 5: (Color online) Coarse grained density of states, $\nu(\varepsilon)$, for the Hamiltonian $H(1, 0.2)$ with $N = 16$. The coarse graining width is $W = 10\delta$, where δ is the mean level spacing. Shown is the result after coarse graining (red circles), and the result after an additional running average (blue line).

$$\chi_W(\varepsilon) = \frac{e^{-\varepsilon^2/(2W^2)}}{\sqrt{2\pi W^2}} \,. \tag{26}$$

In the paper we have used $W = 10\delta$ for $N = 16$, where δ is the mean level spacing with an additional running average. The results of these procedures for the density of states are shown in Fig. 5. The same procedure is performed for the canonical ensemble. As a check that this is working correctly one must compare operator averages found with these coarse grained distributions and with the exact ones. Note that whilst a coarse graining over a wider energy range (*e.g.* $W = 50\delta$) will give the same result for the density of states as in Fig. 5, it does not give accurate results for the canonical ensemble. For the microcanonical ensemble one simply broadens the delta-function around the initial energy $E = \langle\Psi_0|H|\Psi_0\rangle$,

$$\Gamma_{\rm mic}(\varepsilon) \approx \theta(\varepsilon - E + W/2) - \theta(\varepsilon - E - W/2) \,, \tag{27}$$

where, again, $\theta(\varepsilon)$ is the Heaviside function.

In principle one could also attempt this procedure on the diagonal ensemble. However for the system sizes we are able to consider we find that it is not possible to smoothen the diagonal ensemble and, at the same time, retain accurate averages for physical quantities.

The integrable case

Non-equilibrium dynamics in integrable systems and the question of the appropriate statistical ensemble to describe the long-time limit have been intensely studied in recent years.[21, 40, 42–44] In our paper we only briefly touched upon this issue by explaining how the splitting into a local and a non-local part can be generalized to the integrable case. While for non-interacting systems the additional local conservation laws simply become the occupation numbers of the diagonal modes they are quite complicated in the interacting case [28] and a detailed study is beyond the scope of this letter. Here we simply want to look for indications of integrability when taking only the Hamiltonian itself into the local part, as in Eq. (9) of the paper, thus ignoring all other locally conserved quantities. In Fig. 6 we present the same data as in Fig. 2 of the paper but for the integrable quench with $|\Psi_0(5, 0)\rangle$ and $H(1/2, 0)$. The scaling behavior is now quite different. $\Delta_{\rm rel}$ for the correlation function $\langle \boldsymbol{S}_i \boldsymbol{S}_{i+2}\rangle$, in particular, shows an upturn for the largest system sizes indicating that the canonical ensemble does not describe the system at long times. To understand the scaling of the different correlation functions in more detail, an analysis of the overlap of each operator O with the local conserved quantities would be required.



FIG. 6: (Color online) Longer-range correlations for the integrable quench. Shown is the scaling of the relative deviation between ensemble averages $\Delta_{\rm rel}$ for $|i - j| = 1, 2, 3, 4, 5$ (from bottom to top).

Finally, we present in Fig. 7 the analogue of Fig. 3 of the paper for the integrable case. One of the obvious differences is that the fluctuations in O_{nn} for the case $O = \boldsymbol{S}_i \boldsymbol{S}_{i+1}$ are substantially larger than in the non-integrable case, see Fig. 7(b1). In $\bar{O}_{nn}$, shown in Fig. 7(b2) and (c2), the largest fluctuations are of similar magnitude as in the non-integrable case shown in the paper, however, substantial fluctuations persist to much higher energies. This could possibly be related to the different energy level distributions. While level repulsion leads to a Wigner-Dyson distribution for a generic model, each state in an integrable model is uniquely characterized by the quantum numbers of the local conserved quantities so that states can cross. This leads to a Poissonian distribution and might be responsible for the different structure of fluctuations in Fig. 7(b2) and (c2). The fact that we are missing local conservation laws when using the canonical ensemble for an integrable system would, of course, be immediately obvious if we choose



FIG. 7: (Color online) Integrable quench for $N = 16$. Shown are: Shown are: (a) the initial, microcanonical and canonical energy distribution functions, and results for $O = \boldsymbol{S}_i\boldsymbol{S}_{i+1}$ in (b) and $O = \boldsymbol{S}_i\boldsymbol{S}_{i+4}$ in (c). (b1) and (c1) show O_{nn} while (b2) and (c2) show $\bar{O}_{nn}$.

one of the additional local conserved operators as our observable whose expectation value would be time independent and thus characterized by the initial microstate.

[1] L. D. Landau and E. M. Lifshitz, *Statistical Physics* (Butterworth-Heinemann, Oxford, 1980).
[2] G. D. Birkhoff, Proc. Natl. Ac. Sci. **17**, 656 (1931).
[3] J. von Neumann, Proc. Natl. Ac. Sci. **18**, 70 (1932).
[4] G. Morandi, F. Napoli, and E. Ercolessi, *Statistical Mechanics: An intermediate course* (World Scientific, Singapore, 2001).
[5] M. Tabor, *Chaos and integrability in nonlinear dynamics: an introduction* (Wiley, New York, 1989).
[6] J. M. Deutsch, Phys. Rev. A **43**, 2046 (1991).
[7] M. Srednicki, Phys. Rev. E **50**, 888 (1994).
[8] M. Rigol, V. Dunjko, and M. Olshanii, Nature **452**, 854 (2008).
[9] P. Calabrese, F. H. L. Essler, and M. Fagotti, Phys. Rev. Lett. **106**, 227203 (2011).
[10] N. Sedlmayr, J. Ren, F. Gebhard, and J. Sirker, Phys. Rev. Lett. **110**, 100406 (2013).
[11] F. Gebhard, K. zu Münster, J. Ren, N. Sedlmayr, J. Sirker, and B. Ziebarth, Ann. Phys. (Berlin) **524**, 286 (2012).
[12] T. Kinoshita, T. Wenger, and D. S. Weiss, Nature **440**, 900 (2006).
[13] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, Nat. Phys. **8**, 325 (2012).
[14] N. Strohmaier, D. Greif, R. Jördens, L. Tarruell, H. Moritz, T. Esslinger, R. Sensarma, D. Pekker, E. Altman, and E. Demler, Phys. Rev. Lett. **104**, 080401 (2010).
[15] S. Hofferberth, I. Lesanovsky, B. Fischer, T. Schumm, and J. Schmiedmayer, Nature **449**, 324 (2007).
[16] T. Enss and J. Sirker, New J. Phys. **14**, 023008 (2012).
[17] S. R. Manmana, S. Wessel, R. M. Noack, and A. Muramatsu, Phys. Rev. Lett. **98**, 210405 (2007).
[18] M. Rigol, A. Muramatsu, and M. Olshanii, Phys. Rev. A **74**, 053616 (2006).
[19] G. Biroli, C. Kollath, and A. M. Läuchli, Phys. Rev. Lett. **105**, 250401 (2010).
[20] C. Kollath, A. M. Läuchli, and E. Altman, Phys. Rev. Lett. **98**, 180601 (2007).
[21] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Phys. Rev. Lett. **98**, 050405 (2007).
[22] E. Canovi, D. Rossini, R. Fazio, G. E. Santoro, and A. Silva, Phys. Rev. B **83**, 094431 (2011).
[23] M. A. Cazalilla, Phys. Rev. Lett. **97**, 156403 (2006).
[24] L. F. Santos, A. Polkovnikov, and M. Rigol, Phys. Rev. Lett. **107**, 040601 (2011).
[25] T. N. Ikeda, Y. Watanabe, and M. Ueda, Phys. Rev. E **84**, 021130 (2011).
[26] J. von Neumann, Z. Phys. **57**, 30 (1929); Eur. Phys. J. H **35**, 201 (2010).
[27] F. H. L. Essler, H. Frahm, F. Göhmann, A. Klümper, and V. E. Korepin, *The One-Dimensional Hubbard model* (Cambridge University Press, Cambridge, 2005).
[28] M. Grabowski and P. Mathieu, Ann. Phys. **243**, 299 (1995).
[29] G. Roux, Phys. Rev. A **81**, 053604 (2010); Phys. Rev. A **82**, 037602 (2010).
[30] T. Prosen, Phys. Rev. Lett. **106**, 217206 (2011).
[31] X. Zotos, F. Naef, and P. Prelovšek, Phys. Rev. B **55**, 11029 (1997).
[32] A. Rosch and N. Andrei, Phys. Rev. Lett. **85**, 1092 (2000).
[33] J. Sirker, R. G. Pereira, and I. Affleck, Phys. Rev. Lett. **103**, 216602 (2009).
[34] J. Sirker, R. G. Pereira, and I. Affleck, Phys. Rev. B **83**, 035115 (2011).
[35] P. Mazur, Physica **43**, 533 (1969).
[36] M. Suzuki, Physica **51**, 277 (1971).
[37] N. P. Konstantinidis, Phys. Rev. B **72**, 064453 (2005).
[38] P. Barmettler, M. Punk, V. Gritsev, E. Demler, and E. Altman, Phys. Rev. Lett. **102**, 130603 (2009).
[39] P. Barmettler, M. Punk, V. Gritsev, E. Demler, and E. Altman, New J. Phys. **12**, 055017 (2010).
[40] M. Fagotti and F. H. L. Essler, arXiv:1302.6944 (2013).
[41] M. Rigol, Phys. Rev. B **80**, 053607 (2009).
[42] L. F. Santos and M. Rigol, Phys. Rev. E **81**, 036206 (2010).
[43] L. F. Santos and M. Rigol, Phys. Rev. E **82**, 031130 (2010).
[44] J.-S. Caux and R. M. Konik, Phys. Rev. Lett. **109**, 175301 (2012).
[45] In the case of degeneracies, Eq. (2) remains valid if we choose the eigenbasis in which O is diagonal in the degenerate subspaces.